



ELSEVIER

Contents lists available at ScienceDirect

The Journal of Arthroplasty

journal homepage: www.arthroplastyjournal.org

AAHKS Symposium

Innovations in Joint Preservation Procedures for the Dysplastic Hip “The Periacetabular Osteotomy”

Cecilia Pascual-Garrido, MD ^a, Michael D. Harris, PhD ^{b, c}, John C. Clohisy, MD ^{a, *}

^a Adult Reconstruction and Adolescent and Young Adult Hip Service, Department of Orthopaedic Surgery, Washington University School of Medicine, St Louis, Missouri

^b Program in Physical Therapy, Washington University School of Medicine, St Louis, Missouri

^c Department of Orthopaedic Surgery, Washington University School of Medicine, St Louis, Missouri

ARTICLE INFO

Article history:

Received 2 February 2017

Accepted 5 February 2017

Available online xxx

Keywords:

periacetabular osteotomy

hip dysplasia

hip arthroscopy

femoral deformities

developmental dysplasia of the hip

ABSTRACT

The Bernese periacetabular osteotomy is an effective treatment for symptomatic developmental dysplasia in the prearthritic young adult hip. Refinements in the periacetabular osteotomy technique and perioperative management have markedly improved the clinical outcomes and recovery in these patients. We will review the clinical presentation of acetabular dysplasia, indications for surgery, perioperative management, and contemporary refinements in technique including refined acetabular reduction, adjunctive hip arthroscopy, femoral head-neck osteochondroplasty, femoral procedures, and rapid recovery protocols. In well-selected patients, this reconstructive osteotomy should be considered safe and effective in alleviating pain and improving hip function in patients with symptomatic acetabular dysplasia.

© 2017 Elsevier Inc. All rights reserved.

Developmental dysplasia of the hip (DDH) is a common condition associated with pain and functional limitations [1,2]. DDH is also a known risk factor for secondary osteoarthritis (OA), being found in an estimated 20%–40% of secondary OA cases and increasing a patient's risk for OA by 4.3-fold [3,4]. The complex structural deformities of the dysplastic hip can involve both the acetabulum and proximal femur. Acetabular dysplasia is commonly the dominant deformity and is characterized by insufficient femoral head coverage and superolateral inclination of the articular surface. This deformity may result in instability of the joint with overload of the acetabular rim, leading to labral and cartilage damage, and ultimately OA [5,6].

A variety of hip preservation procedures have been developed and proposed for the treatment of symptomatic DDH [7–12]. In 1988, Ganz et al [8] introduced the Bernese periacetabular osteotomy (PAO) for acetabular reorientation. This procedure is

performed through 1 incision, maintains posterior column integrity, preserves the acetabular blood supply, enables versatile multiplanar acetabular reorientation, and provides reliable healing. However, this technique involves a substantial learning curve. Major complications have been reported, especially in the surgeon learning curve phase [13]. Recent innovations in PAO surgery and other joint preservation procedures have had a tremendous positive impact on clinical outcomes of this surgery in the treatment of symptomatic, acetabular dysplasia [14–16].

Here, we present specific protocols which we and others have developed and implemented with a multidisciplinary approach to PAO surgery. These refined protocols have substantially improved the surgical care and postoperative recovery of patients treated with the PAO. The approach includes a consistent plan for pain management, anticoagulation, and active recovery, as well as contemporary refinements to the PAO surgical procedure.

Indications for PAO

The presenting symptoms in patients with symptomatic, acetabular dysplasia can be variable, and radiographic analysis can be challenging, especially in individuals with mild acetabular deformities. Most patients with hip dysplasia present with an insidious onset of activity-related pain localized to the groin and/or the lateral aspect of the hip. Patients normally report

One or more of the authors of this paper have disclosed potential or pertinent conflicts of interest, which may include receipt of payment, either direct or indirect, institutional support, or association with an entity in the biomedical field which may be perceived to have potential conflict of interest with this work. For full disclosure statements refer to <http://dx.doi.org/10.1016/j.arth.2017.02.015>.

* Reprint requests: John C. Clohisy, MD, Adult Reconstruction and Hip Preservation, Department of Orthopaedic Surgery, Washington University School of Medicine, Campus Box 8233, 660 South Euclid Avenue, St Louis, MO 63110.

<http://dx.doi.org/10.1016/j.arth.2017.02.015>

0883-5403/© 2017 Elsevier Inc. All rights reserved.

Table 1
PAO Perioperative Pain Management Protocol.

Preoperative	Intraoperative	Postoperative
(a) Day before surgery: Naprosyn 500 mg PO × 2 tablets	(a) General anesthesia	PACU: (a) Oxycodone 5 mg 1-2 tablets PO × 1
(b) Scopolamine patch (if history of motion sickness or nausea after previous surgery) × 72 h	(b) Time of incision: Tylenol 1 gm IV	(b) Zofran 4 mg IV × 1
	(c) Time of skin closure: Toradol 30 mg IVP	(c) IV Narcotic medication for breakthrough pain as needed
		Discharge medications:
		(a) Naprosyn 500 mg PO bid × 2 wk postoperatively
		(b) Percocet 5/325 1-2 tablets PO 4-6 h PRN—script from hospital
		(c) Hydrocodone/APAP (Norco) 5/325 mg 1-2 tablets PO q 4-6 h PRN after Percocet.

PAO, periacetabular osteotomy; PO, per os; IV, intravenous; PACU, post-anesthesia care unit; PRN, pro re nata; APAP, Acetaminophen; IVP, intravenous push.

moderate-to-severe pain daily and are physiologically young and otherwise healthy. Physical examination findings normally reveal a limp with a positive Trendelenburg sign and a positive impingement test [6].

Acetabular dysplasia pathomorphology is best confirmed using anteroposterior and lateral pelvic radiographs. The modified Dunn view (with hip flexed at an angle of 45° and abducted 20°) is used to detect asphericity of the femoral head and/or head-neck junction deformities, which are known to be common in the dysplastic hip [17–19]. Specifically, patients with a lateral center-edge angle <20°, anterior center-edge angle <18°, and acetabular index >10 are classified as dysplastic [20]. The false-profile view allows further examination of anterior coverage of the femur, confirming dysplasia in those patients with an anterior center-edge angle <18°. False-profile view also helps rule out joint space narrowing in those patients with an anterior subluxation of the femoral head. Additional findings may include rim fractures and fovea alta. Lower surgical threshold should be indicated in patients with DDH and associated laxity, excessive femoral valgus, and/or femur anteversion. For patients with confirmed symptomatic acetabular dysplasia, PAO is indicated for those who are symptomatic, active, and have preserved articular cartilage (Tonnis grade 0-1) after closure of the physis [14,16].

Pain, Anticoagulation, and Blood Loss Management

A preoperative, perioperative, and postoperative pain management protocol has been established to reduce pain before surgery, eliminate the need for an epidural, minimize postoperative nausea

and vomiting, and manage clotting risks. Preoperatively, patients are prescribed nonsteroidal anti-inflammatories (eg, Naprosyn, Roche Laboratories, Indianapolis, IN) for at least 24 hours before surgery (Table 1).

Antinausea medication (eg, scopolamine patch) is also prescribed for 72 hours before surgery to reduce postoperative nausea and vomiting. Intraoperatively, local infiltration anesthesia has been proven to be more effective at controlling pain 12 hours after surgery in comparison with intravenous patient-controlled analgesia [21]. Our protocol consists of general anesthesia with specific timing of intraoperative analgesia followed by patient-controlled analgesia for 12-24 hours. Postoperatively, all patients are managed with intravenous pain medication and progress to oral pain medication as tolerated (Table 1). Multimodality pain management has been a substantial improvement, allowing our patients to have general anesthesia with no epidural. Without the need for an epidural and with minimized postoperative nausea and vomiting, normal discharge time has been expedited to 1-2 days postoperatively.

All patients are indicated to take enteric-coated aspirin 325 mg twice a day for 6 weeks postoperatively for anticoagulation. Intraoperatively, a mobile pneumatic compression device is used on the contralateral leg. In addition, all patients receive tranexamic acid (TXA). TXA inhibits clot breakdown and significantly reduces perioperative blood loss and blood transfusion [22]. Patients with normal renal function normally receive 1 g of TXA, intravenously infused over 10 minutes before skin incision and 1 g infused over 10 minutes at the start of the wound closure [23]. Close monitoring of postoperative hematocrit and hemoglobin is performed in all patients. Cell saver could also be used as an autologous blood recovery system during the procedure.




Fig. 1. Note the minimal incision for periacetabular osteotomy (PAO). (A) Marking pen is used to mark ASIS and iliac crest. Incision is located slightly lateral to the iliac crest. Distally, the incision will be extended further if an additional osteochondroplasty is needed. (B) Open wound after incision is performed. ASIS, anterior superior iliac spine.




Fig. 2. Anteroposterior (AP) pelvis showing bilateral severe hip dysplasia. (A) Note the severe anterolateral undercoverage of the femoral head, the presence of a valgus femur, and the existence of a fatigue “rim fracture” or os acetabuli. (B) At 2 years after PAO, note the excellent correction with preservation of the joint space, good femoral head coverage, and good acetabular inclination and hip center translated medially (low head index extrusion).

Active Recovery

Active recovery for PAO patients includes ambulation with a walker 4–6 hours after operation. Patients are instructed to stay within 30% of full weight bearing for 4 weeks. If the rectus femoris was detached and repaired, to protect the muscle during healing, no active hip flexion exercises or straight leg raises are allowed. However, if the PAO was “rectus sparing” (vast majority of cases), the patient may perform gentle active hip flexion exercises. A continuous passive movement machine is used in all patients for 6 h/d (within comfort level) for 4 weeks with hip flexion to 60°. Patients are not allowed to drive until they are fully weight bearing with active control of the surgical extremity, normally between 6 and 8 weeks. It is also recommended that patients use a raised toilet seat to prevent hip flexion >90°. All patients receive home health physical therapy during the first 4 weeks postoperatively. Patients are then referred to outpatient physical therapy for a total of 20 weeks. Athletic patients are allowed to jog and start impact activities at 3 months. Return to full sport is progressive with a target of full sport at 4 months.

Contemporary Refinements to the PAO Surgical Technique

Advantages of the Bernese PAO originally described by Ganz et al [8], over previous osteotomy techniques, continue to be realized. This review is not intended to have a comprehensive surgical technique section, and as such, the reader is referred to another more comprehensive review on surgical techniques [24].

We perform the PAO with the patient supine on a radiolucent table through a modification of the Smith-Peterson approach (Fig. 1) [25]. Variations of the skin incision and surgical approach exist and are also effective in performing the procedure.

The procedure consists of 4 orthogonal periacetabular cuts that are technically reproducible and allow for substantial multidirectional correction of the mobilized acetabulum. Acetabular reduction should be performed under intraoperative fluoroscopy. Femoral head coverage, acetabular inclination, congruency (joint space), acetabular version, and medial translation of the femoral head should be reestablished (Fig. 2) [20,26,27]. The acetabular blood supply is preserved, which promotes reliable bone healing and minimizes incidents of avascular necrosis of the acetabular fragment [24,28]. Preservation of the posterior column facilitates pelvic and acetabular fragment stability after correction and screw fixation [8].

Contemporary refinements to the surgical technique have focused mostly on improved perioperative management, assessment of acetabular fragment reduction and deformity correction, and treating other hip deformities concomitantly. Additional surgical refinements involve muscle-sparing approaches, with preservation of the abductor muscles (gluteus maximus and medius) and maintenance of the rectus femoris attachment, to improve postoperative recovery [27,29–33]. Concomitant corrective surgeries often involve repair of intraarticular soft tissue (labrum repair and cartilage restoration), femoral osteochondroplasty, decompression of the anterior inferior iliac spine (AIIS), and proximal femoral osteotomy (PFO) to correct femoral dysplasia or severe femoral head deformities.

Intraarticular pathologic changes are a common finding in patients with symptomatic DDH [29]. For addressing and treating intraarticular pathology, arthroscopy can provide suitable visualization and access to address labral, chondral, and ligamentum teres injuries, using a less-invasive approach than open arthrotomy (Fig. 3) [30,34,35]. Still, there is little information about the benefits or risks of performing hip arthroscopy concomitantly with




Fig. 3. Adjunctive hip arthroscopy in a 38-year-old female with left hip dysplasia. (A) Note the left hip with undercoverage of the left femoral head, (B) hip scope confirmed a torn labrum and full-thickness chondral defect, (C) labrum repair and microfracture was performed during arthroscopy, and (D) PAO was performed, immediately after hip arthroscopy, reestablishing good acetabular inclination and lateral coverage of the femoral head.



Fig. 4. Patient with right acetabular dysplasia and associated femoral head-neck offset deformity. Labrum repair, chondroplasty, and microfracture were performed during hip arthroscopy before PAO. Osteochondroplasty of the femoral head-neck junction was performed via anterior arthrotomy at the time of PAO. (A) Note the AP pelvis with right hip dysplasia, (B) Dunn view confirms a cam lesion, (C) arthroscopic microfracture and labrum repair, and (D) femoral head-neck osteochondroplasty is evident.

the PAO as standard of care for the treatment of acetabular dysplasia, and the practice remains controversial [30]. A recently published study, which retrospectively compared PAO alone vs a combined arthroscopy/PAO for the treatment of DDH, suggested better clinical results in those patients with the combined approach [30]. Based on personal preference, some surgeons may perform hip arthroscopy in conjunction with PAO, especially if the patient reports mechanical symptoms, such as catching and popping, but there is a need for a well-conducted, randomized, controlled trial in patients with DDH to confirm the efficacy of this approach. In our practice, we perform hip arthroscopy immediately before PAO, to treat the central compartment, only if the patient reports mechanical symptoms (eg, impingement) and if labral-chondral dissociation has been confirmed by magnetic resonance imaging.

Abnormalities of the femoral head and neck are also common in patients with DDH, and secondary femoroacetabular impingement (FAI) is a frequent mechanism of failure after PAO [17,18,36,37]. Osteochondroplasty is used to alleviate FAI caused by deformities at the femoral head and head-neck junction. Osteochondroplasty can be performed either arthroscopically or through open arthrotomy (Fig. 4) during the PAO procedure by minimal extension of the Smith-Peterson approach distally. Both arthroscopic and open osteochondroplasty have been demonstrated to be safe and effective for managing femoral head-neck deformities [17,38,39]. At our institution, we normally perform an open osteochondroplasty after PAO. Even in the absence of preoperative FAI, it is critical to assess range of motion after acetabular correction during a PAO. If limited range of motion is detected after acetabular correction (limited flexion and internal rotation), open osteochondroplasty is indicated.

Subspine syndrome can be a source of extra-articular impingement in patients with DDH. In these cases, terminal hip flexion results in impingement of the caudal prominence (ie, most distal extension) of the AIIS against the femoral neck and/or anterosuperior greater trochanter [40]. Radiographs and a computed tomography scan are necessary to confirm this pathology and indicate subspine decompression after acetabular fragment correction. Decompression of a prominent AIIS results in improved

hip motion and hip function [40,41]. Subspine decompression can be performed arthroscopically or during PAO procedure through a minimal extension of the modification of the Smith-Peterson approach distally (Fig. 5).

In addition, for patients with severe femoral deformities, such as sequelae of Legg-Calve-Perthes disease or “Perthes-like deformities” or coxa valga with anteverted femurs, an adjunctive PFO may also be required to optimize correction, joint stability, and congruency [7,42–44]. The decision to add a PFO may be made before, during, or after performing PAO [45]. We contemplate adjunctive PFO in 2 clinical situations. First, in the severely dysplastic hip with coxa valga (femoral neck-shaft angle $>140^\circ$) and persistent instability and/or suboptimal congruency with a persistent high head extrusion index after the PAO. Normally, the second main indicator for a PFO is a varus deformity or “Perthes-like” abnormality of the proximal femur. These hips may have a combination of structural and clinical impingement problems, including an aspheric femoral head, short, wide femoral neck, high greater trochanter, limited abduction motion, and leg length discrepancy. Iatrogenic varus deformities are a relatively common indication for concurrent valgus-producing PFO. In these last patients, the PFO is performed first in association with relative femoral neck lengthening, distalization of the trochanter, and valgus osteotomy [46]. The clinical results for patients undergoing concomitant PAO and PFO to address severe hip deformities are comparable with those for isolated PAO [7]. For an in-depth description of surgical correction for patients with DDH and complex femoral deformities, the reader is referred to more comprehensive review articles [7].

Clinical Results

Clinical outcomes after PAO have shown improvement in pain relief and hip function compared with presurgical baselines [18,36,47–51]. However, there is a substantial risk of major complications after PAO, especially during the anticipated surgeon learning curve associated with this procedure [13,52]. Regarding survivorship, recently published literature suggests rates of




Fig. 5. Subspine decompression. (A) Prominent AIIS is evident during open arthrotomy after PAO was performed, (B) open subspine decompression was performed using a burr, (C) arthroscopic view of a prominent AIIS, and (D) arthroscopic decompression and associated labrum repair was performed. AIIS, anterior inferior iliac spine.

Table 2

Clinical Outcomes After PAO at 10 y of Follow-Up or More.

Author	Hips (n)	Age (y)	Follow-Up Years	Survivorship
Wells J; CORR 2017 [18]	121	27 (10–45)	18	85% at 10 y, 80% at 15 y, and 74% at 18 y
Lerch TD; CORR 2016 [49]	75	29 (13–56)	30	29% at 30 y
Matheny T; JBJS 2010 [53]	135	26.7	10	76% at 10 y

CORR, Clinical Orthopaedics and Related Research; JBJS, The Journal of Bone & Joint Surgery; PAO, periacetabular osteotomy.

75%–85% at 10 years and 61% at 20 years follow-up [18,50,53]. When considering long-term outcomes (30 years), only one-third of the hips have shown no progression to OA (Table 2) [49]. However, current long-term outcome data originated from procedures performed in the 1980s when PAO approaches were newly developed and did not have the benefits of the contemporary refinements discussed previously. Most factors that have been associated with progression to OA at 30 years after PAO include the following: older age (>40 years), high preoperative Tonnis grade, decreased preoperative internal rotation, and malcorrection of the acetabular fragment including acetabular overcoverage, or acetabular retroversion [49]. Current post-PAO patient-reported outcome data from the multicenter, prospective Academic Network of Conservative Hip Outcomes Research (ANCHOR) group include 391 hips with a minimum 2-year follow-up and show statistical improvements in clinical score in 93% of patients with a failure rate of 0.8%. Clinical outcomes analysis demonstrated important clinical improvements in pain, function, quality of life, overall health, and activity level. Major complications (classified as modified Clavien-Dindo grades III–IV) were reported in 7% patients including transient nerve palsy, pulmonary emboli, and deep venous thrombosis [16].

Conclusions

Treatment of DDH has advanced remarkably in the past 10 years. Refined patient selection, adequate perioperative management, improved pain management, and contemporary surgical approaches, including the treatment of combined intra-articular pathology and associated impingement, have substantially impacted the outcomes and recovery of patients treated with the PAO. Continued investigation is needed to further refine hip preservation strategies for the dysplastic hip. Identification of predictors of treatment success, failure, and disease modification will further enhance our ability to effectively treat prearthritic and early arthritic dysplastic hip disease.

References

- [1] Klauke K, Durnin CW, Ganz R. The acetabular rim syndrome. A clinical presentation of dysplasia of the hip. *J Bone Joint Surg Br* 1991;73:423–9.
- [2] Gala L, Clohisy JC, Beaulé PE. Hip dysplasia in the young adult. *J Bone Joint Surg Am* 2016;98:63–73.
- [3] Reijman M, Hazes JM, Pols HA, Koes BW, Bierma-Zeinstra SM. Acetabular dysplasia predicts incident osteoarthritis of the hip: the Rotterdam study. *Arthritis Rheum* 2005;52:787–93.
- [4] Aronson J. Osteoarthritis of the young adult hip: etiology and treatment. *Instr Course Lect* 1986;35:119–28.
- [5] Henak CR, Abraham CL, Anderson AE, Maas SA, Ellis BJ, Peters CL, et al. Patient-specific analysis of cartilage and labrum mechanics in human hips with acetabular dysplasia. *Osteoarthritis Cartilage* 2014;22:210–7.
- [6] Nepple JJ, Clohisy JC. The dysplastic and unstable hip: a responsible balance of arthroscopic and open approaches. *Sports Med Arthrosc* 2015;23:180–6.
- [7] Clohisy JC, St John LC, Nunley RM, Schutz AL, Schoenecker PL. Combined periacetabular and femoral osteotomies for severe hip deformities. *Clin Orthop Relat Res* 2009;467:2221–7.

- [8] Ganz R, Klauke K, Vinh TS, Mast JW. A new periacetabular osteotomy for the treatment of hip dysplasias: technique and preliminary results. *Clin Orthop Relat Res* 1988;2004:3–8.
- [9] Ninomiya S, Tagawa H. Rotational acetabular osteotomy for the dysplastic hip. *J Bone Joint Surg Am* 1984;66:430–6.
- [10] Salter RB. The classic. Innominate osteotomy in the treatment of congenital dislocation and subluxation of the hip by Robert B. Salter. *J Bone Joint Surg (Brit)* 43B:3:518, 1961. *Clin Orthop Relat Res* 1978:2–14.
- [11] Steel HH. Triple osteotomy of the innominate bone. *J Bone Joint Surg Am* 1973;55:343–50.
- [12] Tonnis D. Surgical treatment of congenital dislocation of the hip. *Clin Orthop Relat Res* 1990:33–40.
- [13] Novais EN, Carry PM, Kestel LA, Ketterman B, Brusalis CM, Sankar WN. Does surgeon experience impact the risk of complications after Bernese periacetabular osteotomy? *Clin Orthop Relat Res* 2016.
- [14] Coobs BR, Xiong A, Clohisy JC. Contemporary concepts in the young adult hip patient: periacetabular osteotomy for hip dysplasia. *J Arthroplasty* 2015;30:1105–8.
- [15] Leunig M, Ganz R. The evolution and concepts of joint-preserving surgery of the hip. *Bone Joint J* 2014;96-B:5–18.
- [16] Clohisy JC, Ackerman J, Baca G, Baty J, Beaulé PE, Kim YJ, et al. Patient-reported outcomes of periacetabular osteotomy from the Prospective ANCHOR Cohort Study. *J Bone Joint Surg Am* 2017;99:33–41.
- [17] Nassif NA, Schoenecker PL, Thorsness R, Clohisy JC. Periacetabular osteotomy and combined femoral head-neck junction osteochondroplasty: a minimum two-year follow-up cohort study. *J Bone Joint Surg Am* 2012;94:1959–66.
- [18] Wells J, Millis M, Kim YJ, Bulat E, Miller P, Matheny T. Survivorship of the Bernese periacetabular osteotomy: what factors are associated with long-term failure? *Clin Orthop Relat Res* 2017;475:396–405.
- [19] Dunn DM. Anteversion of the neck of the femur; a method of measurement. *J Bone Joint Surg Br* 1952;34-B:181–6.
- [20] Clohisy JC, Carlisle JC, Beaulé PE, Kim YJ, Trousdale RT, Sierra RJ, et al. A systematic approach to the plain radiographic evaluation of the young adult hip. *J Bone Joint Surg Am* 2008;90(Suppl 4):47–66.
- [21] Novais EN, Kestel L, Carry PM, Sink E, Strupp K. Local infiltration analgesia compared with epidural and intravenous PCA after surgical hip dislocation for the treatment of femoroacetabular impingement in adolescents. *J Pediatr Orthop* 2016 [Epub ahead of print]. PMID: 26840273.
- [22] Mont MA, Jacobs JJ. AAOS clinical practice guideline: preventing venous thromboembolic disease in patients undergoing elective hip and knee arthroplasty. *J Am Acad Orthop Surg* 2011;19:777–8.
- [23] Alshryda S, Mason J, Vaghela M, Sarda P, Nargol A, Maheswaran S. Topical (intra-articular) tranexamic acid reduces blood loss and transfusion rates following total knee replacement: a randomized controlled trial (TRANX-K). *J Bone Joint Surg Am* 2013;95:1961–8.
- [24] Clohisy JC, Barrett SE, Gordon JE, Delgado ED, Schoenecker PL. Periacetabular osteotomy in the treatment of severe acetabular dysplasia. Surgical technique. *J Bone Joint Surg Am* 2006;88(Suppl 1 Pt 1):65–83.
- [25] Smith-Petersen MN. Approach to and exposure of the hip joint for mold arthroplasty. *J Bone Joint Surg Am* 1949;31A:40–6.
- [26] Lehmann CL, Nepple JJ, Baca G, Schoenecker PL, Clohisy JC. Do fluoroscopy and postoperative radiographs correlate for periacetabular osteotomy corrections? *Clin Orthop Relat Res* 2012;470:3508–14.
- [27] Wylie JD, Ross JA, Erickson JA, Anderson MB, Peters CL. Operative fluoroscopic correction is reliable and correlates with postoperative radiographic correction in periacetabular osteotomy. *Clin Orthop Relat Res* 2017;475:1100–6. <http://dx.doi.org/10.1007/s11999-016-5071-1>.
- [28] Beck M, Leunig M, Ellis T, Sledge JB, Ganz R. The acetabular blood supply: implications for periacetabular osteotomies. *Surg Radiol Anat* 2003;25:361–7.
- [29] Ross JR, Zaltz I, Nepple JJ, Schoenecker PL, Clohisy JC. Arthroscopic disease classification and interventions as an adjunct in the treatment of acetabular dysplasia. *Am J Sports Med* 2011;39(Suppl):72S–8S.
- [30] Ricciardi BF, Mayer SW, Fields KG, Wentzel C, Kelly BT, Sink EL. Patient characteristics and early functional outcomes of combined arthroscopic labral refixation and periacetabular osteotomy for symptomatic acetabular dysplasia. *Am J Sports Med* 2016;44:2518–25.
- [31] Clohisy JC, Nunley RM, Carlisle JC, Schoenecker PL. Incidence and characteristics of femoral deformities in the dysplastic hip. *Clin Orthop Relat Res* 2009;467:128–34.
- [32] Novais EN, Kim YJ, Carry PM, Millis MB. The Bernese periacetabular osteotomy: is transection of the rectus femoris tendon essential? *Clin Orthop Relat Res* 2014;472:3142–9.
- [33] Peters CL, Erickson JA, Anderson MB, Anderson LA. Preservation of the rectus femoris origin during periacetabular osteotomy does not compromise acetabular reorientation. *Clin Orthop Relat Res* 2015;473:608–14.
- [34] Domb BG, LaReau JM, Hammarstedt JE, Gupta A, Stake CE, Redmond JM. Concomitant hip arthroscopy and periacetabular osteotomy. *Arthroscopy* 2015;31:2199–206.
- [35] Nepple JJ, Carlisle JC, Nunley RM, Clohisy JC. Clinical and radiographic predictors of intra-articular hip disease in arthroscopy. *Am J Sports Med* 2011;39:296–303.
- [36] Steppacher SD, Tannast M, Ganz R, Siebenrock KA. Mean 20-year followup of Bernese periacetabular osteotomy. *Clin Orthop Relat Res* 2008;466:1633–44.
- [37] Myers SR, Eijer H, Ganz R. Anterior femoroacetabular impingement after periacetabular osteotomy. *Clin Orthop Relat Res* 1999:93–9.

- [38] Nwachukwu BU, Rebolledo BJ, McCormick F, Rosas S, Harris JD, Kelly BT. Arthroscopic versus open treatment of femoroacetabular impingement: a systematic review of medium- to long-term outcomes. *Am J Sports Med* 2016;44:1062–8.
- [39] Goronzy J, Franken L, Hartmann A, Thielemann F, Postler A, Paulus T, et al. What are the results of surgical treatment of hip dysplasia with concomitant cam deformity? *Clin Orthop Relat Res* 2017;475:1128–37. <http://dx.doi.org/10.1007/s11999-016-5054-2>.
- [40] Hetsroni I, Poultsides L, Bedi A, Larson CM, Kelly BT. Anterior inferior iliac spine morphology correlates with hip range of motion: a classification system and dynamic model. *Clin Orthop Relat Res* 2013;471:2497–503.
- [41] Hetsroni I, Larson CM, Dela Torre K, Zbeda RM, Magennis E, Kelly BT. Anterior inferior iliac spine deformity as an extra-articular source for hip impingement: a series of 10 patients treated with arthroscopic decompression. *Arthroscopy* 2012;28:1644–53.
- [42] Albers CE, Steppacher SD, Schwab JM, Tannast M, Siebenrock KA. Relative femoral neck lengthening improves pain and hip function in proximal femoral deformities with a high-riding trochanter. *Clin Orthop Relat Res* 2015;473:1378–87.
- [43] Ganz R, Horowitz K, Leunig M. Algorithm for femoral and periacetabular osteotomies in complex hip deformities. *Clin Orthop Relat Res* 2010;468:3168–80.
- [44] Clohisy JC, Nepple JJ, Ross JR, Pashos G, Schoenecker PL. Does surgical hip dislocation and periacetabular osteotomy improve pain in patients with Perthes-like deformities and acetabular dysplasia? *Clin Orthop Relat Res* 2015;473:1370–7.
- [45] Hersche O, Casillas M, Ganz R. Indications for intertrochanteric osteotomy after periacetabular osteotomy for adult hip dysplasia. *Clin Orthop Relat Res* 1998;19–26.
- [46] Clohisy JC, Ross JR, North JD, Nepple JJ, Schoenecker PL. What are the factors associated with acetabular correction in Perthes-like hip deformities? *Clin Orthop Relat Res* 2012;470:3439–45.
- [47] Clohisy JC, Barrett SE, Gordon JE, Delgado ED, Schoenecker PL. Periacetabular osteotomy for the treatment of severe acetabular dysplasia. *J Bone Joint Surg Am* 2005;87:254–9.
- [48] Hartig-Andreasen C, Troelsen A, Thillemann TM, Søballe K. What factors predict failure 4 to 12 years after periacetabular osteotomy? *Clin Orthop Relat Res* 2012;470:2978–87.
- [49] Lerch TD, Steppacher SD, Liechti EF, Tannast M, Siebenrock KA. One-third of hips after periacetabular osteotomy survive 30 years with good clinical results, no progression of arthritis, or conversion to THA. *Clin Orthop Relat Res* 2017;475:1154–68. <http://dx.doi.org/10.1007/s11999-016-5169-5>.
- [50] Troelsen A, Elmengaard B, Soballe K. Medium-term outcome of periacetabular osteotomy and predictors of conversion to total hip replacement. *J Bone Joint Surg Am* 2009;91:2169–79.
- [51] Matheny T, Zaltz I, Kim YJ, Schoenecker P, Millis M, Podeszwa D, et al. Activity level and severity of dysplasia predict age at Bernese periacetabular osteotomy for symptomatic hip dysplasia. *J Bone Joint Surg Am* 2016;98:665–71.
- [52] Zaltz I, Baca G, Kim YJ, Schoenecker P, Trousdale R, Sierra R, et al. Complications associated with the periacetabular osteotomy: a prospective multicenter study. *J Bone Joint Surg Am* 2014;96:1967–74.
- [53] Matheny T, Kim YJ, Zurakowski D, Matero C, Millis M. Intermediate to long-term results following the Bernese periacetabular osteotomy and predictors of clinical outcome: surgical technique. *J Bone Joint Surg Am* 2010;92(Suppl 1 Pt 2):115–29.